arXiv:1012.3674v2 [math.CV] 17 Dec 2010

Another extension of the disc algebra*

V. Nestoridis† and N. Papadatos

Department of Mathematics, University of Athens,
Panepistemiopolis, 157 84 Athens, Greece.

Abstract

We identify the complex plane $\mathbb{C}$ with the open unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$ by the homeomorphism $\mathbb{C} \ni z \mapsto \frac{z}{1+|z|} \in D$. This leads to a compactification $\overline{\mathbb{C}}$ of $\mathbb{C}$, homeomorphic to $\overline{D} = \{z \in \mathbb{C} : |z| \leqslant 1\}$. The Euclidean metric on $\overline{D}$ induces a metric d on $\overline{\mathbb{C}}$. We identify all uniform limits of polynomials on $\overline{D}$ with respect to the metric d. The class of the above limits is an extension of the disc algebra and it is denoted by $\overline{A}(D)$. We study properties of the elements of $\overline{A}(D)$ and topological properties of the class $\overline{A}(D)$ endowed with its natural topology. The class $\overline{A}(D)$ is different and, from the geometric point of view, richer than the class $\widetilde{A}(D)$ introduced in [6], [7], on the basis of the chordal metric χ.

MSC: Primary 30J99, secondary 46A99, 30E10.

Key words and phrases: Disc Algebra; Mergelyan's Theorem; Polynomial Approximation.

1 Introduction

The uniform limits of the polynomials on the closed unit disc $\overline{D} = \{z \in \mathbb{C} : |z| \leqslant 1\}$ with respect to the usual Euclidean metric on $\mathbb{C}$ are exactly all functions $f : \overline{D} \to \mathbb{C}$, continuous on $\overline{D}$ and holomorphic in $D = \{z \in \mathbb{C} : |z| < 1\}$. The class of the above functions is the disc algebra $A(D)$. In [6], [7], the Euclidean metric on $\mathbb{C}$ is replaced by the chordal metric χ on the one-point compactification $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ of $\mathbb{C}$. The set of uniform limits of polynomials, with respect to χ, is an extension of $A(D)$ and it is denoted by $\widetilde{A}(D)$. It contains the constant function equal to ∞ and the following functions: $f : \overline{D} \to \widetilde{\mathbb{C}}$ continuous, such that $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic. The class $\widetilde{A}(D)$ remains the same if we replace the chordal metric χ by any other metric on $\widetilde{\mathbb{C}}$ generating the same topology – the reason is that any two such metrics are uniformly equivalent, because $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ is compact. In this sense, the set

*Work partially supported by the University of Athens' Research fund under Grants 70/4/4923 and 70/4/5637.

†Corresponding author. e-mail address: vnestor@math.uoa.gr

of χ-uniform polynomial limits is an *invariant set*, i.e., independent of the specific metric one chooses for generating it.

Instead of the one-point compactification $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ we shall consider another, more rich from the geometric view-point, compactification, as follows: We identify the complex plane $\mathbb{C}$ with the open unit disc D by the homeomorphism

$$\mathbb{C} \ni z \mapsto \frac{z}{1+|z|} \in D.$$

The natural compactification of D is $\overline{D}$. This leads to a compactification $\overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$ of $\mathbb{C}$, where the set of infinite points $\mathbb{C}^\infty$ is homeomorphic to the unit circle $T = \partial D = \{z \in \mathbb{C} : |z| = 1\}$; more precisely we write $\mathbb{C}^\infty = \{\infty e^{i\theta} : \theta \in \mathbb{R}\}$. The usual Euclidean metric on $\overline{D}$ induces the metric d on $\overline{\mathbb{C}}$, defined as follows:

$$d(z_1, z_2) = \begin{cases} \left| \dfrac{z_1}{1+|z_1|} - \dfrac{z_2}{1+|z_2|} \right|, & \text{for } z_1, z_2 \in \mathbb{C}, \\ \left| \dfrac{z_1}{1+|z_1|} - e^{i\theta} \right|, & \text{for } z_1 \in \mathbb{C},\ z_2 = \infty e^{i\theta}\ (\theta \in \mathbb{R}), \\ \left| e^{i\theta_1} - e^{i\theta_2} \right|, & \text{for } z_1 = \infty e^{i\theta_1},\ z_2 = \infty e^{i\theta_2}\ (\theta_1, \theta_2 \in \mathbb{R}). \end{cases} \tag{1.1}$$

We shall investigate the uniform limits (on $\overline{D}$) of the polynomials, with respect to the metric d. The class of the limit functions is another extension of the disc algebra, different from $\widetilde{A}(D)$, which is denoted by $\overline{A}(D)$. It will be shown that it contains exactly two types of functions. The first type, the *finite type*, consists of all continuous functions $f : \overline{D} \to \overline{\mathbb{C}}$ such that $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic. The second type, the *infinite type*, consists of all functions $f : \overline{D} \to \mathbb{C}^\infty \subset \overline{\mathbb{C}}$ of the form $f(z) = \infty e^{i\theta(z)}$ where the function $\theta : \overline{D} \to \mathbb{R}$ is continuous on $\overline{D}$ and harmonic in D.

To highlight the difference between $\widetilde{A}(D)$ and $\overline{A}(D)$ we mention that, e.g., the function $f(z) = \frac{1}{1-z}$, with $f(1) = \infty$, belongs to $\widetilde{A}(D)$ but not to $\overline{A}(D)$. An example of an element of $\overline{A}(D)$ of the finite type, not belonging to $A(D)$, is given by $f(z) = \log \frac{1}{1-z}$ (here $f(1) = +\infty$ corresponds to the infinite element $\infty e^{i\theta} \in \mathbb{C}^\infty \subset \overline{\mathbb{C}}$ with $\theta = 0$).

Furthermore, we shall investigate some properties of the elements of $\overline{A}(D)$ and some of its topological properties when it is endowed with its natural metric. Finally, we shall consider uniform approximation with respect to the metric d on other compact sets, different from $\overline{D}$. Certainly, due to the compactness of $\overline{\mathbb{C}}$, all the above results remain valid if we replace d by any equivalent metric on $\overline{\mathbb{C}}$.

Several open questions are naturally posed and new directions of investigation are indicated. In particular, any result on $A(D)$ and any approximation result with respect to the usual Euclidean metric is worth to be examined in $\overline{A}(D)$ and with respect to d, respectively.

2 Preliminaries

Let $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the one-point compactification of the complex plane $\mathbb{C}$, endowed by the chordal metric χ, defined by $\chi(z_1, z_2) = \frac{|z_1 - z_2|}{\sqrt{1+|z_1|^2}\sqrt{1+|z_2|^2}}$ for $z_1, z_2 \in \mathbb{C}$, $\chi(z_1, \infty) = \frac{1}{\sqrt{1+|z_1|^2}}$ for $z_1 \in \mathbb{C}$ and, certainly, $\chi(\infty, \infty) = 0$.

We can also define another compactification $\overline{\mathbb{C}}$ of $\mathbb{C}$ with infinitely many points of infinity, as follows: First consider the homomorphism $G : \mathbb{C} \to \{w \in \mathbb{C} : |w| < 1\} = D$ given by $G(z) = \frac{z}{1+|z|}$. Since $\overline{D}$ is a compactification of D, it induces a compactification $\overline{\mathbb{C}}$ of $\mathbb{C}$. The set $\mathbb{C}^\infty$, which consists of all points of infinity, is homeomorphic with the circle $T = \partial D = \{w \in \mathbb{C} : |w| = 1\}$; thus, $\overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$. Every element of $\mathbb{C}^\infty$ is determined by a unimodular complex number $e^{i\theta}$, $\theta \in \mathbb{R}$, and we shall denote the corresponding element of $\mathbb{C}^\infty \subset \overline{\mathbb{C}}$ by $\infty e^{i\theta}$. In particular, this compactification contains the usual two points of infinity of the real numbers ($\pm\infty$), in the sense that $+\infty$ corresponds to $\theta = 0$ while $-\infty$ is related to $\theta = \pi$. The Euclidean metric on $\overline{D}$ induces the metric d on $\overline{\mathbb{C}}$, which is defined in (1.1). The definition of the metric d can be simplified if we extend $G(z) = \frac{z}{1+|z|}$, defined for $z \in \mathbb{C}$, to the points of infinity $z = \infty e^{i\theta} \in \mathbb{C}^\infty$ ($\theta \in \mathbb{R}$), as $G(z) = G(\infty e^{i\theta}) = e^{i\theta}$. Then, $d(z_1, z_2) = |G(z_1) - G(z_2)|$ for all $z_1, z_2 \in \overline{\mathbb{C}}$.

Both metrics χ and d induce the usual topology on $\mathbb{C}$. For if $W \subset \mathbb{C}$ is compact, then χ, d and the Euclidean metric (all restricted on W) are two by two uniformly equivalent. Thus, if S is any set and $f, f_n : S \to W$ ($n = 1, 2, \ldots$) are functions, then the uniform convergence on S, $f_n \to f$, as $n \to +\infty$, with respect to any one of these metrics, implies uniform convergence with respect to the other two. Further, we have the following

Lemma 2.1 If S is any set and $f, f_n : S \to \mathbb{C}$ ($n = 1, 2, \ldots$) are functions, then the uniform convergence $f_n \to f$, as $n \to \infty$, with respect to the usual Euclidean metric on $\mathbb{C}$, implies the uniform convergence $f_n \to f$, as $n \to \infty$, with respect to the metric d.

Proof: The result is implied by the fact that for all $z_1, z_2 \in \mathbb{C}$,

$$d(z_1, z_2) \leqslant |z_1 - z_2|. \tag{2.1}$$

This inequality can be proved as follows. Write

$$d(z_1, z_2) = \frac{|z_1 - z_2 + w|}{1 + |z_1| + |z_2| + |z_1 z_2|}, \quad \text{where} \quad w = z_1|z_2| - z_2|z_1|,$$

and observe that $w = \dfrac{|z_1| + |z_2|}{2}(z_1 - z_2) - \dfrac{z_1 + z_2}{2}(|z_1| - |z_2|)$. It follows that

$$|z_1 - z_2 + w| \leqslant \left(1 + \frac{|z_1| + |z_2|}{2}\right)|z_1 - z_2| + \frac{|z_1| + |z_2|}{2}\Big||z_1| - |z_2|\Big| \leqslant (1 + |z_1| + |z_2|)|z_1 - z_2|,$$

and thus,

$$d(z_1, z_2) \leqslant \left(\frac{1 + |z_1| + |z_2|}{1 + |z_1| + |z_2| + |z_1 z_2|} \right) |z_1 - z_2| \leqslant |z_1 - z_2|.$$

□

Consider now the map $\Phi : (\overline{\mathbb{C}}, d) \to (\widetilde{\mathbb{C}}, \chi)$, defined for all $z \in \overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$ by

$$\Phi(z) = \begin{cases} z, & \text{if } z \in \mathbb{C}, \\ \infty, & \text{if } z = \infty e^{i\theta} \in \mathbb{C}^\infty \ (\theta \in \mathbb{R}). \end{cases} \tag{2.2}$$

One can easily see that Φ is continuous and, therefore, uniformly continuous, because $\overline{\mathbb{C}}$ is compact. This immediately implies the following

Lemma 2.2 If S is any set and $f, f_n : S \to \overline{\mathbb{C}}$ $(n = 1, 2, \ldots)$ are functions, then the uniform convergence (on S) $f_n \to f$, as $n \to +\infty$, with respect to the metric d, implies the uniform convergence $\Phi \circ f_n \to \Phi \circ f$, as $n \to +\infty$, with respect to the metric χ.

Remark 2.1 Alternatively, one can verify Lemma 2.2 by making use of the easily proved inequality $\chi(\Phi(z_1), \Phi(z_2)) \leqslant 2d(z_1, z_2)$, which is valid for all $z_1, z_2 \in \overline{\mathbb{C}}$.

Corollary 2.1 If S is any set and $f, f_n : S \to \mathbb{C}$ $(n = 1, 2, \ldots)$ are functions, then the uniform convergence (on S) $f_n \to f$, as $n \to +\infty$, with respect to the usual Euclidean metric on $\mathbb{C}$ implies the uniform convergence $f_n \to f$, as $n \to +\infty$, with respect to the metric χ.

Proof: It suffices to combine Lemmas 2.1 and 2.2, or to observe the trivial inequality $\chi(z_1, z_2) \leqslant |z_1 - z_2|$, $z_1, z_2 \in \mathbb{C}$. □

Another useful fact that will be used in the sequel is the following; its simple proof is omitted.

Lemma 2.3 Let $R > 0$ be a positive real number and define the map $\Phi_R : \overline{\mathbb{C}} \to \mathbb{C}$ by

$$\Phi_R(z) = \begin{cases} z, & \text{if } z \in \mathbb{C} \ \text{ and } \ |z| < R, \\ R\dfrac{z}{|z|}, & \text{if } z \in \mathbb{C} \ \text{ and } \ |z| \geqslant R, \\ Re^{i\theta}, & \text{if } z = \infty e^{i\theta} \in \mathbb{C}^\infty \ (\theta \in \mathbb{R}). \end{cases}$$

Then, Φ_R is (uniformly) continuous.

3 The definition

In this section we consider the closed unit disc $S = \overline{D} = \{z \in \mathbb{C} : |z| \leqslant 1\}$ and we identify the set of uniform limits on $\overline{D}$ of polynomial functions with respect to the metric d.

Suppose f_n, $n = 1, 2, \ldots$, is a sequence of complex polynomials. Let $f : \overline{D} \to \overline{\mathbb{C}}$ be a function, where $\overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$, endowed with the metric d, is the compactification of $\mathbb{C}$ introduced previously. We assume that the sequence f_n, $n = 1, 2, \ldots$, converges uniformly on $\overline{D}$ towards f with respect to the metric d. Since polynomial functions are continuous and uniform convergence preserves continuity, it follows that the limiting function $f : \overline{D} \to \overline{\mathbb{C}}$ has to be continuous. Furthermore, according to Lemma 2.2, the sequence $\Phi \circ f_n = f_n$, $n = 1, 2, \ldots$, converges uniformly (with resect to χ) to the function $\Phi \circ f : \overline{D} \to \widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. Here Φ is the map defined by (2.2) and χ is the chordal metric on $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ (see section 2). It follows that the function $\Phi \circ f$ belongs to the class $\widetilde{A}(D)$ introduced in [6], [7]. Thus, according to the definition of $\widetilde{A}(D)$, the function $\Phi \circ f$ can be of the following two types:

The first type contains the holomorphic functions $\Phi \circ f : D \to \mathbb{C}$, such that for every boundary point $\zeta \in \partial D = T = \{w \in \mathbb{C} : |w| = 1\}$, the limit

$$\lim_{z \to \zeta,\ z \in D} \Phi(f(z))$$

exists in $\widetilde{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$. In this case we conclude that the continuous function $f : \overline{D} \to \overline{\mathbb{C}}$ satisfies $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic. This is the first type, the *finite type* of $\overline{A}(D)$.

The second type of elements of $\widetilde{A}(D)$ is the constant function $\Phi \circ f \equiv \infty$; this means that for every $z \in \overline{D}$, the value $f(z)$ is a point of infinity of $\overline{\mathbb{C}}$. In this case the continuous function $f : \overline{D} \to \overline{\mathbb{C}}$ satisfies $f(\overline{D}) \subset \mathbb{C}^\infty$. Thus, f is of the form $f(z) = \infty e^{i\theta(z)}$ for some function $\theta : \overline{D} \to \mathbb{R}$. Observe that the continuity of the function $f(z) = \infty e^{i\theta(z)}$, with respect to the metric d, is equivalent to the continuity of the function $e^{i\theta(z)}$, with respect to the usual Euclidean metric. Thus, the function

$$\overline{D} \ni z \mapsto e^{i\theta(z)} \in T = \partial D$$

has to be continuous. Since $\overline{D}$ is simply connected, it follows from Theorem 5.1 of [5] (see page 128) that the real function $\overline{D} \ni z \mapsto \theta(z) \in \mathbb{R}$ can be chosen to be continuous. Now we shall show that any continuous version of the function $\overline{D} \ni z \mapsto \theta(z) \in \mathbb{R}$ is, in fact, harmonic in the open unit disc D.

Indeed, the uniform convergence to the $\mathbb{C}^\infty$-valued function f (with respect to the metric d) shows that $|f_n(z)| \to +\infty$, as $n \to +\infty$, uniformly on $\overline{D}$. Thus, $|f_n(z)| \geqslant 1$ for all $z \in \overline{D}$ and for all $n \geqslant n_0$. Considering in D a branch of $\log f_n$ we conclude that $\mathrm{Arg} f_n(z) = \mathrm{Im}[\log f_n(z)]$ is a harmonic function in D. We also have $\frac{f_n(z)}{|f_n(z)|} \to e^{i\theta(z)}$, uniformly on $\overline{D}$ (with respect to the usual Euclidean metric). Thus, the same holds on $D \subset \overline{D}$. It follows that $e^{i[\theta(z) - \mathrm{Arg} f_n(z)]} \to 1$, as $n \to +\infty$, uniformly on D. Thus, there exists $n_1 \geqslant n_0$, such that for every natural number $n \geqslant n_1$, there exists an

integer $k_n = k_n(z) \in \mathbb{Z}$ such that the function $w_n(z) := \theta(z) - \mathrm{Arg} f_n(z) - 2k_n(z)\pi$ takes values in $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ for all $z \in D$. Since $e^{iw_n(z)} = \frac{e^{i\theta(z)}}{e^{i\mathrm{Arg} f_n(z)}}$ is continuous in D and $w_n(z) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$, it follows that the function $D \ni z \mapsto w_n(z) \in \mathbb{R}$ is continuous $(n \geqslant n_1)$. Therefore, writing $k_n(z) = \frac{1}{2\pi}(\theta(z) - w_n(z) - \mathrm{Arg} f_n(z))$ we see that the function $D \ni z \mapsto k_n(z) \in \mathbb{Z}$ is also continuous, and hence, constant. Thus, for $z \in D$ we may write $k_n(z) \equiv k_n \in \mathbb{Z}$, independent of $z \in D$ (for $n \geqslant n_1$). Now, the uniform convergence $e^{iw_n(z)} \to 1$, as $n \to +\infty$ (with respect to the Euclidean metric) and the fact that $w_n(z) \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ for all $z \in D$ and for all $n \geqslant n_1$, imply that $w_n(z) \to 0$, uniformly on D (with respect to the Euclidean metric). Equivalently, $2k_n\pi + \mathrm{Arg} f_n(z) \to \theta(z)$, as $n \to +\infty$, uniformly on D. Since the functions $2k_n\pi + \mathrm{Arg} f_n(z)$, $n \geqslant n_1$, are harmonic in D, it follows that the function $\theta(z)$, being a uniform limit of harmonic functions, is also harmonic in D.

Therefore, we have shown that the second type of the limiting functions are continuous functions $f : \overline{D} \to \mathbb{C}^\infty \subset \overline{\mathbb{C}}$, of the form $f(z) = \infty e^{i\theta(z)}$, where the function $\overline{D} \ni z \mapsto \theta(z) \in \mathbb{R}$ can be chosen to be continuous on $\overline{D}$ and harmonic in D. The above $\mathbb{C}^\infty$-valued functions generate the *infinite type* of $\overline{A}(D)$.

Conversely, we shall show that each continuous function $f : \overline{D} \to \overline{\mathbb{C}}$ of the previous two types is, indeed, the uniform limit (with respect to the metric d) of a sequence of polynomials $f_n : \overline{D} \to \mathbb{C}$. The details are as follows:

Suppose first that f is of the finite type. That is, $f : \overline{D} \to \overline{\mathbb{C}}$ is continuous (with respect to the metric d), $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic. Since $\overline{D}$ is compact, it follows that f is uniformly continuous. Therefore, for a given $\epsilon > 0$, we can find a real number $r = r(\epsilon)$, $0 < r < 1$, such that $d(f(z), f(rz)) < \frac{\epsilon}{2}$ for all $z \in \overline{D}$. Since the function $f : D \to \mathbb{C}$ is holomorphic, a partial sum $P(w)$ of the Taylor development of $f(w)$ satisfies $|f(w) - P(w)| < \frac{\epsilon}{2}$ for all $w : |w| \leqslant r$. Thus, $|f(rz) - P(rz)| < \frac{\epsilon}{2}$ for all $z \in \overline{D}$. Using inequality (2.1) we see that $d(f(rz), P(rz)) < \frac{\epsilon}{2}$ for all $z \in \overline{D}$. Now, the triangle inequality implies that $d(f(z), Q(z)) < \epsilon$, $z \in \overline{D}$, where $Q = Q_\epsilon$ is the polynomial defined by $Q(z) = P(rz)$. Setting $\epsilon = 1/n$ we conclude that the sequence of polynomials $f_n = Q_{1/n}$, $n = 1, 2, \ldots$, approximates f uniformly on $\overline{D}$, with respect to d.

Finally suppose that f is of the infinite type. That is, assume that $f : \overline{D} \to \mathbb{C}^\infty \subset \overline{\mathbb{C}}$ is continuous (with respect to d) and of the form $f(z) = \infty e^{i\theta(z)}$, $z \in \overline{D}$, where the real-valued function $\theta(z)$, $z \in \overline{D}$, is continuous on $\overline{D}$ and harmonic in D. Let $\epsilon > 0$. Since the function $\overline{D} \ni z \mapsto \theta(z) \in \mathbb{R}$ is uniformly continuous, there exists $r = r(\epsilon)$, with $0 < r < 1$, such that $|\theta(z) - \theta(rz)| < \frac{\epsilon}{3}$ for all $z \in \overline{D}$. Therefore, from $d(f(z), f(rz)) = |e^{i\theta(z)} - e^{i\theta(rz)}| \leqslant |\theta(z) - \theta(rz)|$ we conclude that $d(f(z), f(rz)) < \frac{\epsilon}{3}$ for all $z \in \overline{D}$. Since the function $D \ni z \mapsto \theta(z) \in \mathbb{R}$ is harmonic in the simply connected domain D, we can find a holomorphic function $g \in \mathcal{H}(D)$ such that $\mathrm{Im} g(z) = \theta(z)$, $z \in D$. Setting $\delta = \min\{e^{\mathrm{Re} g(w)} : |w| \leqslant r\}$ we see that $\delta > 0$. Thus, for any $n \in \{1, 2, \ldots\}$ and any $w \in \mathbb{C}$ with $|w| \leqslant r$, we have

$$d(ne^{g(w)}, f(w)) = \frac{1}{1 + ne^{\mathrm{Re} g(w)}} \leqslant \frac{1}{1 + \delta n}.$$

It follows that we can fix a large enough $n = n(\epsilon)$ so that $d(ne^{g(rz)}, f(rz)) < \frac{\epsilon}{3}$ for all $z \in \overline{D}$. On the other hand, since the function $ne^{g(w)}$ is holomorphic in D, it can be approximated, uniformly over $\overline{D}_r = \{w \in \mathbb{C} : |w| \leqslant r\}$, by a partial sum of its Taylor expansion, with respect to the Euclidean metric. Therefore, there exists a polynomial $P(w)$ such that $|ne^{g(w)} - P(w)| < \frac{\epsilon}{3}$ for all $w : |w| \leqslant r$. Using the inequality (2.1) (cf. Lemma 2.1) we conclude that $d(ne^{g(rz)}, P(rz)) < \frac{\epsilon}{3}$ for all $z \in \overline{D}$. By considering the polynomial $Q = Q_\epsilon$, defined by $Q(z) = P(rz)$, we conclude that for all $z \in \overline{D}$,

$$d(f(z), Q_\epsilon(z)) \leqslant d(f(z), f(rz)) + d(f(rz), ne^{g(rz)}) + d(ne^{g(rz)}, P(rz)) < \epsilon.$$

It follows that the sequence of polynomials $f_n = Q_{1/n}$, $n = 1, 2, \ldots$, approximates f uniformly on $\overline{D}$, with respect to d.

Therefore, we naturally arrived at the following definition.

Definition 3.1 Let $\mathbb{C}^\infty = \{\infty e^{i\theta} : \theta \in \mathbb{R}\}$ and set $\overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$, endowed by the metric d defined in (1.1). Let also $D = \{z \in \mathbb{C} : |z| < 1\}$ and $\overline{D} = \{z \in \mathbb{C} : |z| \leqslant 1\}$ be, as usually, the open and the closed unit disc, respectively. We denote by $\overline{A}(D)$ the class of continuous functions $f : \overline{D} \to \overline{\mathbb{C}}$ of the following two types.

(a) The finite type: It contains the continuous functions $f : \overline{D} \to \overline{\mathbb{C}}$ such that $f(D) \subset \mathbb{C}$ and $f_{|D}$ is holomorphic in D.

(b) The infinite type: It contains the continuous functions $f : \overline{D} \to \mathbb{C}^\infty \subset \overline{\mathbb{C}}$ of the form $f(z) = \infty e^{i\theta(z)}$, $z \in \overline{D}$, where the real-valued function $\theta : \overline{D} \to \mathbb{R}$ is harmonic in D and continuous (with respect to the usual Euclidean metric) on $\overline{D}$.

Thus, we have shown the following

Theorem 3.1 Let $\overline{D}$ be the closed unit disc. The set of uniform limits (on $\overline{D}$) of the complex-valued polynomials – with respect to the metric d, defined on $\overline{\mathbb{C}} = \mathbb{C} \cup \mathbb{C}^\infty$ by (1.1) – coincides with the class $\overline{A}(D)$.

Obviously $\overline{A}(D)$ contains the disc algebra $A(D) = \{f : \overline{D} \to \mathbb{C}$, continuous on $\overline{D}$, holomorphic in $D\}$. It is an extension of $A(D)$, essentially different than the extension $\widetilde{A}(D)$ obtained in [6], [7], using the chordal metric χ. Indeed, it is easily seen that the function $f(z) = \frac{1}{1-z}$, $|z| < 1$, is neither a restriction in D of any element of $A(D)$, nor a restriction in D of any element of $\overline{A}(D)$. However, $f(z)$ can be extended to $\overline{D}$ in an obvious manner, so that the resulting $\widetilde{\mathbb{C}}$-valued function is χ-continuous and forms an element of $\widetilde{A}(D)$. Loosely speaking, $f(z) = \frac{1}{1-z}$ belongs to $\widetilde{A}(D)$ but not to $\overline{A}(D)$.

Note that any conformal mapping of D onto an open strip (or half-strip) belongs to $\overline{A}(D)$ and not to $A(D)$. More generally, it is also true that for any nonzero complex numbers c_k, $k = 1, 2, \ldots, n$, the function

$$f(z) = \sum_{k=1}^{n} c_k \log \frac{1}{e^{i\theta_k} - z}, \quad (\text{where } 0 \leqslant \theta_1 < \theta_2 < \cdots < \theta_n < 2\pi)$$

is an element of $\overline{A}(D)$, and not of $A(D)$. In fact, the above examples belong to the finite type of $\overline{A}(D)$. It is easy to show that if f belongs to $\overline{A}(D)$ (and is of finite type) then $\Phi \circ f$ belongs to $\widetilde{A}(D)$ – see (2.2), Lemma 2.2 and Remark 2.1. On the other hand the converse does not hold, as we can see by the example $f(z) = \frac{1}{1-z}$; obviously, this function cannot be written as $f = \Phi \circ g$ for some $g \in \overline{A}(D)$.

Some simple examples of elements of $\overline{A}(D)$ of the infinite type can be constructed as follows: Consider a polynomial P not vanishing at any point of $\overline{D}$. Then, the sequence $nP(z)$, $n = 1, 2, \ldots$, converges uniformly on $\overline{D}$, with resect to the metric d, to the function $f(z) = \infty e^{i\mathrm{Arg}P(z)}$ which, certainly, belongs to the infinite type of $\overline{A}(D)$. In particular, taking $P(z) = (2+z)^k$ with large enough $k \in \mathbb{N}$ (in fact, $k = 6$ suffices), we see that the image of the limiting function f covers the whole $\mathbb{C}^\infty$.

4 Some properties of the elements of $\overline{A}(D)$

Let $f \in \overline{A}(D)$. Then $\Phi \circ f \in \widetilde{A}(D)$ and, applying Proposition 3.1 of [6], [7], we obtain the following.

Proposition 4.1 Let $T = \partial D = \{\zeta \in \mathbb{C} : |\zeta| = 1\}$ be the unit circle and assume that $f \in \overline{A}(D)$.
(a) If for some $c \in \mathbb{C}$ the set $\{\zeta \in T : f(\zeta) = c\}$ has positive Lebesgue measure then f is constant.
(b) If the set $\{\zeta \in T : f(\zeta) \in \mathbb{C}^\infty\}$ has positive Lebesgue measure then f is of the infinite type.

Remark 4.1 If $f(z) = \infty e^{i\theta(z)}$ belongs to the infinite type of $\overline{A}(D)$ then $\theta(z)$ can be constant on a subarc of T with strictly positive length without being constant on $\overline{D}$. In fact, any continuous function $T \ni \zeta \mapsto \theta(\zeta) \in \mathbb{R}$ has a unique extension $\overline{D} \ni z \mapsto \theta(z) \in \mathbb{R}$ which is continuous on $\overline{D}$ and harmonic in D; it defines a unique $f \in \overline{A}(D)$ of infinite type.

Proposition 4.2 Let K be a compact subset of T having Lebesgue measure zero. Then any continuous (with respect to the metric d) function $\phi : K \to \mathbb{C}^\infty$ is the restriction of some $f \in \overline{A}(D)$, of finite type, such that $f^{-1}(\mathbb{C}^\infty) = K$.

Proof: There exists a function $g \in A(D)$ such that $g(z) = 1$ on K and $|g(z)| < 1$ on $\overline{D} \setminus K$ (see [2], p. 81). Also, since $\phi(K) \subset \mathbb{C}^\infty$, we can write $\phi(\zeta) = \infty e^{i\theta(\zeta)}$, $\zeta \in K$, for some real-valued function $\theta : K \to \mathbb{R}$. Since the function ϕ is continuous (on K) with respect to d, the function $e^{i\theta(\zeta)}$ is continuous (on K) with respect to the usual Euclidean metric. It follows from [2], [1], [8] that there exists a function $h \in A(D)$ such that $h(\zeta) = e^{i\theta(\zeta)}$ for $\zeta \in K$. Now it is easy to verify that the function given by

$$f(z) = \begin{cases} h(z) \log \frac{1}{1-g(z)}, & \text{if } z \in \overline{D} \setminus K, \\ \phi(z), & \text{if } z \in K, \end{cases}$$

has the desired properties. □

If $E \subset T$ is compact with positive Lebesgue measure, then E is not a compact of interpolation for $\overline{A}(D)$. That is, there exists a continuous function $\eta : E \to \overline{\mathbb{C}}$ which does not have an extension $f \in \overline{A}(D)$. Indeed, let $\zeta_0 \in E$ and let V be an arc with middle point ζ_0 and length less than the one half of the Lebesgue measure of E. We set $\eta \equiv 0$ on $E \setminus V$, $\eta(\zeta_0) = 1$, and we extend η linearly on V. Assume now that for some $f \in \overline{A}(D)$ it is true that $f_{|E} \equiv \eta$. Then $f_{|E\setminus V} \equiv 0$. Since $E \setminus V$ has positive Lebesgue measure, Proposition 4.1(a) implies that $f \equiv 0$, which contradicts $f(\zeta_0) = \eta(\zeta_0) = 1$.

Question 1 If $E \subset T$ is a compact set with Lebesgue measure zero, is it true that E is a compact of interpolation of $\overline{A}(D)$? That is, is it true that every continuous function $\eta : E \to \overline{\mathbb{C}}$ has an extension in $\overline{A}(D)$?
[We refer to [1], [8] for the corresponding result for $A(D)$.]

One can easily see that for any compact set $E \subset T$, every continuous function $\eta : E \to \mathbb{C}^\infty$ has an extension in $\overline{A}(D)$ of the infinite type; this follows from the fact that every continuous function $\theta : E \to \mathbb{R}$ has a continuous extension on $\overline{D}$ which is harmonic in D. Of course, this extension is unique only in the case $E = T$.

Another question is as follows:

Question 2 Characterize the compact sets $E \subset \overline{D}$ having the property that every continuous function $\eta : E \to \overline{\mathbb{C}}$, with $\eta(E \cap D) \subset \mathbb{C}$, has an extension in $\overline{A}(D)$.
[We refer to [3] for the corresponding result for $A(D)$.]

One can also pose questions on the nature of the zero set of a function $f \in \overline{A}(D)$ of the finite type. Also, what can be said about the nature of the set $\{z \in \overline{D} : f(z) = \infty e^{i0}\} = \{z \in \overline{D} : f(z) = +\infty\} = f^{-1}(+\infty)$, when f is of the infinite type? This is related to the zero sets (in $\overline{D}$) of functions $\theta : \overline{D} \to \mathbb{R}$ which are continuous on $\overline{D}$ and harmonic in D.

The maximum principle does not hold in $\overline{A}(D)$. Indeed, consider the polynomials $f(z) = z$ and $g(z) = 2z$, which certainly belong to $A(D) \subset \overline{A}(D)$. Then $d(f(z), g(z)) = \frac{|z|}{(1+|z|)(1+2|z|)}$. For $|z| = 1$ we find $d(f(z), g(z)) = \frac{1}{6} < \frac{1}{3+2\sqrt{2}} = d\left(f\left(\frac{1}{\sqrt{2}}\right), g\left(\frac{1}{\sqrt{2}}\right)\right)$. However, we have the following

Proposition 4.3 Let $f, g \in \overline{A}(D)$ and suppose that $f(\zeta) = g(\zeta)$ for all $\zeta \in T$. Then $f \equiv g$.

Proof: Consider the set $A = \{\zeta \in T : f(\zeta) \in \mathbb{C}^\infty\} = \{\zeta \in T : g(\zeta) \in \mathbb{C}^\infty\}$. If A has positive Lebesgue measure then both f and g are of infinite type. Write $f(z) = \infty e^{i\theta(z)}$ and $g(z) = \infty e^{i\phi(z)}$ where $\theta, \phi : \overline{D} \to \mathbb{R}$ are continuous functions on $\overline{D}$, harmonic in D. Since $f(\zeta) = g(\zeta)$ for all $\zeta \in T$ we conclude that $\theta(\zeta) = \phi(\zeta) + 2k\pi$ for all $\zeta \in T$, where k is an integer independent of $\zeta \in T$. This implies that $\theta(z) = \phi(z) + 2k\pi$ for all $z \in \overline{D}$ and, thus, $f \equiv g$.

Suppose now that A has Lebesgue measure zero. Then f and g are both of finite type. Thus, $f(D) \subset \mathbb{C}$, $g(D) \subset \mathbb{C}$ and both f, g are holomorphic in D. Therefore,

the function $f-g$ is holomorphic in D with zero limits on $T\setminus A$. Since $T\setminus A$ contains a compact set of positive Lebesgue measure, Privalov's Theorem ([4], p. 84) implies $f\equiv g$. This completes the proof. $\square$

Remark 4.2 Assume that $f,g\in\overline{A}(D)$ coincide on a compact set $E\subset T$ with positive Lebesgue measure. If f is of finite type then g is also of finite type and Privalov's Theorem ([4], p. 84) implies $f\equiv g$. If, however, f and g are of infinite type, it may happen $f\neq g$. For example, set $\theta(\zeta)=0$ on $\{\zeta\in T:\zeta=e^{it},\ 0\leqslant t\leqslant\pi\}$ and consider two different continuous (real-valued) extensions θ_1, θ_2 on T. Extending θ_1 and θ_2 on $\overline{D}$, using the Poisson kernel, we find that the functions $f(z)=\infty e^{i\theta_1(z)}$ and $g(z)=\infty e^{i\theta_2(z)}$ belong to $\overline{A}(D)$, coincide on $\{e^{it},0\leqslant t\leqslant\pi\}$ but $f\neq g$.

One more question is the following.

Question 3 Applying Lebesgue dominated convergence Theorem, or by a direct computation, one can see that the function $f(z)=\log\frac{1}{1-z}$ satisfies the mean value property:

$$f(0)=\frac{1}{2\pi}\int_0^{2\pi}f(e^{i\theta})d\theta.$$

Is this the case for all $f\in\overline{A}(D)$, of finite type?

5 Some topological properties of $\overline{A}(D)$

We recall that $A(D)=\{f:\overline{D}\to\mathbb{C}$, continuous on $\overline{D}$ and holomorphic in $D\}$ is a Banach algebra if it is endowed with the usual supremum norm. Furthermore, $\widetilde{A}(D)=\{f:\overline{D}\to\widetilde{\mathbb{C}}=\mathbb{C}\cup\{\infty\}$, continuous on $\overline{D}$, $f(D)\subset\mathbb{C}$, $f_{|D}$ holomorphic$\}\cup\{f:f\equiv\infty\}$ is a complete metric space if it is endowed with the metric $\widetilde{\chi}$ given by (see [6], [7])

$$\widetilde{\chi}(f,g)=\sup_{|z|\leqslant 1}\chi(f(z),g(z)),\quad f,g\in\widetilde{A}(D).$$

Finally, $\overline{A}(D)$ is naturally endowed with the metric

$$\overline{d}(f,g)=\sup_{|z|\leqslant 1}d(f(z),g(z)),\quad f,g\in\overline{A}(D).$$

Proposition 5.1 The metric space $(\overline{A}(D),\overline{d})$ is complete. The disc algebra $A(D)$ is an open and dense subset of $\overline{A}(D)$. The relative topology of $A(D)$ from $\overline{A}(D)$ coincides with the usual topology of $A(D)$.

Proof: Consider the set $\mathcal{B}=\{f:\overline{D}\to\overline{\mathbb{C}}\}$ endowed with the metric

$$\beta(f,g)=\sup_{|z|\leqslant 1}d(f(z),g(z)),\quad f,g\in\mathcal{B}.$$

Since $(\overline{\mathbb{C}},d)$ is complete, it follows that $(\mathcal{B},\beta)$ is complete. According to Theorem 3.1, $\overline{A}(D)$ is the closure in $\mathcal{B}$ of the set of polynomials. Thus, $\overline{A}(D)$ is a closed subset

of the complete metric space $\mathcal{B}$. It follows that $\overline{A}(D)$ is also complete. Let $f \in A(D)$; then $f(\overline{D})$ is a compact subset of $\mathbb{C}$ and, obviously, the compact sets $f(\overline{D})$ and $\mathbb{C}^\infty$ are disjoint. Thus, $\text{dist}(f(\overline{D}), \mathbb{C}^\infty) = \delta > 0$. It is easily seen that if a function $g \in \overline{A}(D)$ satisfies $\overline{d}(f, g) < \delta$ then $g(\overline{D}) \subset \mathbb{C}$ and hence, $g \in A(D)$. Therefore, $A(D)$ is an open subset of $\overline{A}(D)$. It is also dense, because it contains the set of polynomials which is dense, according to Theorem 3.1.

Let $f, f_n \in A(D)$ $(n = 1, 2, \ldots)$ and assume that $f_n \to f$, as $n \to +\infty$, in $A(D)$. From Lemma 2.1 we easily see that $f_n \to f$, as $n \to +\infty$, in $\overline{A}(D)$. Conversely, assume that $f_n \to f$, as $n \to +\infty$, in $\overline{A}(D)$, for some functions $f, f_n \in A(D)$ $(n = 1, 2, \ldots)$. Then, the set $f(\overline{D})$ is a compact subset of $\mathbb{C}$ which is disjoint of the compact set $\mathbb{C}^\infty$, so that $\text{dist}(f(\overline{D}), \mathbb{C}^\infty) = \delta > 0$. Thus, for $n \geqslant n_0$ we have $f_n(\overline{D}) \subset \{w \in \overline{\mathbb{C}} : \text{dist}(w, f(\overline{D})) \leqslant \delta/2\} = E$, say, which is a compact subset of $\mathbb{C}$. On E the usual Euclidean metric and the metric d are uniformly equivalent. It follows that $f_n \to f$, as $n \to +\infty$, in $A(D)$. This completes the proof. $\square$

Consider now the function $\mathbb{F} : \overline{A}(D) \to \widetilde{A}(D)$ defined by $\mathbb{F}(f)(z) = \Phi(f(z))$, i.e.,

$$\overline{A}(D) \ni f \mapsto \mathbb{F}(f) = \Phi \circ f \in \widetilde{A}(D),$$

(see Lemma 2.2 and (2.2) for the definition of the map $\Phi : \overline{\mathbb{C}} \to \widetilde{\mathbb{C}}$). According to Lemma 2.2 the function $\mathbb{F}$ is continuous.

Corollary 5.1 The set $\mathbb{F}(\overline{A}(D))$ is a dense subset of $\widetilde{A}(D)$; in fact it is residual.

Proof: Since $\mathbb{F}(\overline{A}(D))$ contains the set of polynomials, it is dense in $\widetilde{A}(D)$ ([6], [7]). Also, $\mathbb{F}(\overline{A}(D))$ contains $A(D)$ which is open and dense in $\widetilde{A}(D)$ ([6], [7]). Thus, $\mathbb{F}(\overline{A}(D))$ is residual in $\widetilde{A}(D)$. $\square$

Also, it is easily seen that the elements of finite type of $\overline{A}(D)$ form an open dense subset, say $\overline{A}_\circ(D)$, of $\overline{A}(D)$. It follows that the elements of the infinite type of $\overline{A}(D)$ form the closed subset $\overline{A}_\infty(D) = \overline{A}(D) \setminus \overline{A}_\circ(D)$, which is of the first category. Since $\overline{A}_\infty(D)$ is closed, it is a G_δ set.

In the following proposition we use some notation from [10].

Proposition 5.2 Let η be any Hausdorff measure function. The set of all $f \in \overline{A}_\circ(D)$ such that $\Lambda_\eta(E_f) = 0$ is dense and G_δ in $\overline{A}(D)$, where $E_f = \{\zeta \in T : f(\zeta) \notin f(D)\}$.

The proof is similar to the proof of Proposition 4.3 of [6], [7], the only difference being that one has to consider $f^{-1}(\mathbb{C}^\infty)$ in place of $f^{-1}(\infty)$.

Next we define $\mathcal{Y} = \{f \in \overline{A}(D) : f(D) \subset f(T)\} \subset \overline{A}(D)$ and $\mathcal{W} = \{f \in \overline{A}(D) : f(T) = \overline{\mathbb{C}}\} \subset \overline{A}(D)$. Arguments similar to those given in Proposition 4.5 of [6], [7] show that $\mathcal{Y}$ is a non-empty closed subset of $\overline{A}(D)$ of the first category. With a proof similar to the proof of Proposition 4.6 in [6], [7], we can show that $\mathcal{W}$ is also a closed subset of $\overline{A}(D)$ of the first category, but we do not know if $\mathcal{W}$ is non-empty. However, if we assume that every compact set $K \subset T$ with zero Lebesgue measure

is a compact of interpolation for $\overline{A}(D)$, then we can show that $\mathcal{W} \neq \emptyset$. Indeed, let $K \subset T$ be a Cantor-type set with Lebesgue measure zero. It is well known that there exists a continuous surjection $\phi : K \to [0,1]$. Let $\Gamma : [0,1] \to \overline{D}$ be a Peano curve with $\Gamma([0,1]) = \overline{D}$. Finally let $L : \overline{D} \to \overline{\mathbb{C}}$ be a homeomorphism. Then $L \circ \Gamma \circ \phi$ is continuous on K with $(L \circ \Gamma \circ \phi)(K) = \overline{\mathbb{C}}$. Therefore, the assumption that K is a compact of interpolation for $\overline{A}(D)$ implies that there exists an $f \in \overline{A}(D)$ such that $f_{|K} = L \circ \Gamma \circ \phi$. This f belongs to $\mathcal{W}$.

6 Concluding remarks and questions

In the previous sections we considered uniform approximation by polynomials on the compact set $\overline{D}$. However, we can also consider uniform approximation on other compact sets with respect to the metric d. Also, the approximating functions do not necessarily have to be polynomials.

Proposition 6.1 Let $L \subset \mathbb{C}$ be a compact set and let $z_0 \in L^0$. We assume that for every boundary point $\zeta \in \partial L$ the segment $[z_0, \zeta]$ satisfies $[z_0, \zeta] \setminus \{\zeta\} \subset L^0$. Then, the uniform limits, with respect to the metric d, of polynomials on L are exactly the functions $f : L \to \overline{\mathbb{C}}$ of the following two types:

(a) The first type (the finite type) contains the continuous functions $f : L \to \overline{\mathbb{C}}$ with $f(L^0) \subset \mathbb{C}$ such that $f_{|L^0}$ is holomorphic.

(b) The second type (the infinite type) contains the continuous functions $f : L \to \mathbb{C}^\infty$ of the form $f(z) = \infty e^{i\theta(z)}$, where the function $\theta : L \to \mathbb{R}$ can be chosen to be continuous on L and harmonic in L^0.

For the proof we may assume $z_0 = 0$ and imitate the proof for the case $L = \overline{D}$. The difference is that when we approximate f on the compact set rL, $0 < r < 1$, we have to use Runge's Theorem rather than considering the Taylor expansion of f. Since the approximation is uniform on rL with respect to the Euclidean distance on $\mathbb{C}$, Lemma 2.1 shows that it is also uniform with respect to the metric d.

Theorem 6.1 Let $f : T \to \overline{\mathbb{C}}$ be any continuous function. Then, there exists a sequence of trigonometric polynomials converging to f uniformly on T with respect to the metric d.

Proof: Let $\epsilon > 0$. We wish to find a complex-valued trigonometric polynomial $Q = Q_\epsilon$ such that $d(f(\zeta), Q(\zeta)) < \epsilon$ for all $\zeta \in T$. According to Lemma 2.3, for any $R > 0$ the composition $\Phi_R \circ f : T \to \mathbb{C}$ is continuous on T. Fix a large enough $R > 0$ so that $d(f(\zeta), \Phi_R(f(\zeta))) < \frac{\epsilon}{2}$ for all $\zeta \in T$. Now, since $\Phi_R \circ f$ takes only finite complex values, it can be uniformly approximated on T by a trigonometric polynomial Q with respect to the usual Euclidean metric on $\mathbb{C} \cong \mathbb{R}^2$. According to Lemma 2.1, the approximation remains uniform for the metric d. Thus, we have found a trigonometric polynomial Q such that $d(Q(\zeta), \Phi_R(f(\zeta))) < \frac{\epsilon}{2}$ for all $\zeta \in T$. The triangle inequality now yields the desired result. $\square$

We mention here that we do not know what is the set of the uniform limits of polynomials, with respect to the metric d, on a circle.

Let now I be a compact segment in $\mathbb{C}$ or, more generally, a homeomorphic image of the segment $[0,1]$ in $\mathbb{C}$. Then, the uniform limits of the polynomials on I, with respect to the metric d, are exactly all continuous functions $f : I \to \overline{\mathbb{C}}$. The proof is similar to that of Theorem 6.1, with the difference that we make use of the classical Mergelyan's Theorem to approximate $\Phi_R \circ f$ by a (complex) polynomial. This is possible because $I^0 = \emptyset$ and $\mathbb{C} \setminus I$ is connected.

Fix now I to be the compact interval $[-1,1]$. So far, we have seen approximations of $\overline{\mathbb{C}}$-valued functions by complex-valued polynomials. However, when a continuous function $f : [-1,1] \to \overline{\mathbb{C}}$ is $\overline{\mathbb{R}}$-valued, where $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\} \subset \overline{\mathbb{C}}$, it is reasonable to approximate it, if possible, by real-valued polynomials, with respect to the metric d. For the same reasoning, any χ-continuous $\widetilde{\mathbb{R}}$-valued function $f : [-1,1] \to \widetilde{\mathbb{R}} \subset \widetilde{\mathbb{C}}$, where $\widetilde{\mathbb{R}} = \mathbb{R} \cup \{\infty\}$, should be approximated by real-valued polynomials with respect to the metric χ. According to [6], [7], any continuous function $f : [-1,1] \to \widetilde{\mathbb{C}}$, and hence, any continuous function $f : [-1,1] \to \widetilde{\mathbb{R}}$, can be uniformly approximated by *complex-valued* polynomials with respect to the metric χ; however, the approximating polynomials need not be real and, sometimes, they cannot be real. Consider, for example, the χ-continuous function $f : [-1,1] \to \widetilde{\mathbb{R}}$, given by

$$f(x) = \begin{cases} \dfrac{1}{x}, & \text{if } x \in [-1,0) \cup (0,1], \\ \infty, & \text{if } x = 0. \end{cases}$$

Although there exist polynomial approximations for this f, it is easily seen that the approximating polynomials cannot be real-valued – the above function is χ-continuous and not d-continuous. In fact, one can show that a function $f : [-1,1] \to \widetilde{\mathbb{R}}$ can be uniformly approximated by *real-valued* polynomials, with respect to the metric χ, if and only if it is of the form $f = \Phi \circ g$ for some d-continuous function $g : [-1,1] \to \overline{\mathbb{R}}$; here the map $\Phi : \overline{\mathbb{R}} \to \widetilde{\mathbb{R}}$ is the restriction on $\overline{\mathbb{R}} \subset \overline{\mathbb{C}}$ of the map Φ defined in (2.2). Also, it is easy to see that a function $f : [-1,1] \to \overline{\mathbb{R}}$ can be uniformly approximated by *real-valued* polynomials, with respect to the metric d, if and only if it is d-continuous. In other words, uniform real polynomial approximations (with respect to the metric d) can be found for a function $f : [-1,1] \to [-\infty, +\infty]$ if and only if for each $x_0 \in [-1,1]$, $\lim_{x \to x_0} f(x) = f(x_0) \in [-\infty, +\infty]$. If this is true then the same real polynomials approximate $\Phi \circ f$ in the χ-metric. For example, the function $f : [-1,0) \cup (0,1] \to \mathbb{R}$ with $f(x) = \frac{1}{x^2}$ (for $x \in [-1,1]$, $x \neq 0$) can be extended, in an obvious manner, to a d-continuous function on $[-1,1]$ (setting $f(0) = +\infty$) and to a χ-continuous function on $[-1,1]$ (setting $f(0) = \infty$). It follows that this f can be uniformly approximated by real polynomials with respect to the metric d (and, hence, also with respect to χ).

Finally, it is natural to ask about the uniform limits of polynomials on L, with respect to the metric d, when L is a compact subset of $\mathbb{C}$ with connected complement. Specifically, we have the following

Question 4 Let $L \subset \mathbb{C}$ be a compact set with connected complement. Let $f : L \to \overline{\mathbb{C}}$ be a continuous function, such that for every component V of L^0, the following holds: either $f(V) \subset \mathbb{C}$ and $f_{|V}$ is holomorphic, or $f(V) \subset \mathbb{C}^\infty$ and f is of the form $f(z) = \infty e^{i\theta(z)}$ for all $z \in V$, where the real-valued function θ is harmonic in V. Does there exist a sequence of (complex-valued) polynomials converging to f uniformly on L with respect to the metric d?

The existence of such a sequence of polynomials would lead to an extension of the classical Mergelyan's Theorem in the case of the metric d; we refer to [9] for the classical Mergelyan's Theorem.

We notice that the converse is true and the proof is the same as the one given here for the particular case $L = \overline{D}$. Indeed, in the proof of Theorem 3.1 we have only used the fact that D is a simply connected domain; this is the case for every component V of L^0.

Acknowledgement. We thank D. Cheliotis for some helpful comments regarding the metric d.

References

[1] L. Carleson, Representation of continuous functions, *Math. Zeitschr. Bd* 66S. 447–451 (1957).

[2] K. Hoffman, *Banach Spaces of Analytic Functions*, M.I.T., Prentice-Hall, Inc. Englewood Cliffs, N.J.

[3] Val' Der Kastro and M.S. Mel'nikov, On interpolation sets for the algebra $R(X)$, *Izv. Akad. Nauk SSSR*, Ser. Mat. Tom 41 (1997), No 2.

[4] P. Koosis, *Introduction to H_p spaces*, London Math. Soc. Lecture Note Series 40, Cambridge Univ. Press, Cambridge, London, N.Y. New Rochelle, Melbourne, Sydney.

[5] William S. Massey (1965). *A Basic Course in Algebraic Topology*, Sringer-Verlag (1991), New York, Berlin, Heibelberg, London, Paris, Tokyo, Hong Kong, Barcelona.

[6] V. Nestoridis, An extension of the disc algebra, I, submitted.

[7] V. Nestoridis, An extension of the disc algebra, Arxiv: 1009.5364.

[8] W. Rudin, Boundary values of continuous Analytic Functions, *Proc. A.M.S.*, **7**(5), 808–811 (1956).

[9] W. Rudin, *Real and Complex Analysis*, McGraw-Hill, N.Y., St. Louis, San Francisco, Toronto, London, Sydney.

[10] D. Stegenga and K. Stephenson, Generic covering properties for spaces of Analytic Functions, *Pacific J. Math.*, **119**(6), 227–243 (1985).